

METHODS AND RISK FACTORS FOR EXAMINING PATIENTS WITH GOUT

Sharapova Nozima Erkinjonovna

Asian International University, Bukhara, Uzbekistan

Abstract: *Gout is a crystal-deposition disease that results from chronic elevation of uric acid levels above the saturation point for monosodium urate (MSU) crystal formation. Initial presentation is mainly severely painful episodes of peripheral joint synovitis (acute self-limiting 'attacks') but joint damage and deformity, chronic usage-related pain and subcutaneous tophus deposition can eventually develop. The global burden of gout is substantial and seems to be increasing in many parts of the world over the past 50 years. However, methodological differences impair the comparison of gout epidemiology between countries. In this comprehensive Review, data from epidemiological studies from diverse regions of the world are synthesized to depict the geographic variation in gout prevalence and incidence. Key advances in the understanding of factors associated with increased risk of gout are also summarized. The collected data indicate that the distribution of gout is uneven across the globe, with prevalence being highest in Pacific countries. Developed countries tend to have a higher burden of gout than developing countries, and seem to have increasing prevalence and incidence of the disease. Some ethnic groups are particularly susceptible to gout, supporting the importance of genetic predisposition. Socioeconomic and dietary factors, as well as comorbidities and medications that can influence uric acid levels and/or facilitate MSU crystal formation, are also important in determining the risk of developing clinically evident gout.*

Keywords: *Gender; Gout; Risk factors; Systematic review*

Independent risk factors for gout. Obesity is one of the main components associated with gout in the long term . Obesity is a well-established risk factor for

the development of diabetes, dyslipidemia, cardiovascular disease, hypertension, osteoarthritis, and gout. Epidemiological studies have shown that body mass index is independently associated with increased uric acid levels [18, 20]. Many clinical and epidemiological studies have been conducted on the association of hyperuricemia with various cardiovascular diseases. Some authors consider hyperuricemia to be an independent risk factor for the development of cardiovascular disease and an increased risk of mortality in patients . There are many studies in the literature, including epidemiological studies on the combination and interaction of hypertension and hyperuricemia.

Increased blood uric acid levels are considered by many authors as a predictor of the development of hypertension. Studies conducted in sufficiently large cohorts of patients have revealed an increased risk of hypertension even in patients with relatively low blood uric acid levels (300-420 $\mu\text{mol/l}$). Hypertension developing in patients with hyperuricemia and gout can be considered secondary. This is explained by a number of factors: the deposition of uric acid crystals in the interstitial tissues leads to the formation of a giant cell reaction with excessive production of cytokines, followed by fibrosis of the interstitial tissues of the kidneys and the development of tubulointerstitial nephritis. The development of tubulointerstitial kidney damage leads to increased sodium reabsorption in the ascending limb of the loop of Henle and increased sensitivity to natriuretic hormone, leading to water and sodium retention, increased circulating blood volume, total peripheral vascular resistance, activation of the sympathetic division, the nervous system, and ultimately to the development of hypertension. In addition, glomerular hyalinosis and intimal hypertrophy may develop. Another reason for the development of hypertension against the background of hyperuricemia may be a decrease in the response of the renin-angiotensin system to angiotensin II due to a decrease in the sensitivity of angiotensin II receptors in patients .

Left ventricular hypertrophy is common in patients with hypertension and hyperuricemia. Increased left ventricular myocardial mass index in patients with

hypertension and established myocardial hypertrophy is closely related to uricemia [16]. Many studies have shown that the prooxidant effect of SC on lipid oxidation, particularly low-density lipoprotein, is an assessment of oxidative stress in patients with acute myocardial infarction and acute heart failure [3, 14]. The prooxidant effect of SC has also been found to be associated with cardiovascular disease, atherosclerosis, diabetes, lipid peroxidation, and increased endothelial dysfunction products. The effect of SC on endothelial dysfunction is associated with the development of hypertension by the expression of adhesion molecules on the surface of endothelial cells under the influence of the pro-inflammatory cytokines interleukin-1 (IL-1) and tumor necrosis factor. In these patients, impaired nitrogen release from endothelial cells and an increase in C-reactive protein are observed. The relationship between hyperuricemia and AG is still unclear. The presence of hypertension itself may be the cause of the formation of GU. If we take into account that AG is formed as a result of the action of uric acid, AG itself may contribute to the subsequent development of hyperuricemia. This depends on a number of factors. Hypertension causes microvascular damage, which in turn leads to tissue ischemia and the breakdown of ATP into adenine and xanthine, increased production of xanthine oxidase, and, as a result, the formation of purines from nucleosides due to hyperuricemia. Oxygen radicals are used in the xanthine oxidase reaction, which, in turn, activates lipid peroxidation. Thus, timely identification of risk factors is the basis for the prevention of cardiovascular events caused by atherosclerosis. It is of great importance to study renal function and the state of the lipid profile in patients with gout, especially in combination with arterial hypertension.

Methods of diagnosing gout

Despite the clear pathogenesis of the disease and the availability of effective therapy, gout is often not diagnosed or is diagnosed late. According to V.A. Nasonova et al. (2004), gout is diagnosed only in the 8th year of the disease.

For the diagnosis of gout, only the diagnostic criteria of the Wallace S.L. classification (1997) and the classification criteria approved by WHO in 2000 are

currently considered to be the only ones. The criteria consist of three blocks: I (A) - the presence of characteristic crystalline urates in the joint fluid or II (B) - the presence of tophi containing crystalline urates confirmed by chemical or polarizing microscopy or III (C) - the presence of 6 of the following 12 signs.

1. multiple arthritis attacks in the anamnesis;
2. maximum inflammation on the first day;
3. monoarticular nature of arthritis;
4. hyperemia of the skin over the affected joint;
5. swelling and pain in the first big toe joint;
6. unilateral damage to the first big toe joint;
7. unilateral damage to the joints of the arch of the foot;
8. tophi;
9. hyperuricemia;
10. asymmetric swelling of the affected joint;
11. subcortical cysts without erosion on radiographs;
12. Lack of flora in the synovial fluid during synovial fluid culture.

In the absence of a polarizing microscope, the diagnosis of gout can be made according to clinical criteria III (B) (the presence of 6 of the 12 above signs), but in 100% of cases, a reliable diagnosis can be made only by the presence of characteristic crystalline urates in the synovial fluid and / or the presence of a tophi containing crystalline urates confirmed by chemical or polarizing microscopy. The diagnosis of gout seems simple even without microscopy. The nature of acute gouty arthritis is very characteristic: acute pain, reaching the culmination of arthritis within 12 hours, involvement of the first metatarsal phalanx (MTFB) joint, asymmetry. However, the presence of MTFB I is highly sensitive (0.98; 95% confidence interval 0.95–1.02) but has low specificity (0.23; 95% CI 0.10–0.35). Acute monoarthritis, including MTFB I, is also seen in other diseases [12]. In this regard, the European League Against Rheumatism (EULAR) expert panel recommends that physicians perform joint puncture in patients suspected of having gout, as well as in any other undifferentiated arthritis [8].

Several studies have shown that polarizing microscopy has a high specificity for detecting sodium monourate crystals (SMC) in synovial fluid or in tophus, but there is interlaboratory variability. This is likely due to the fact that SMC crystals cannot be detected in all patients with significant gout in synovial fluid samples. This depends on the stage of the disease, allopurinol therapy, and a number of other factors.

Hyperuricemia is the most important risk factor for gout. However, uric acid levels may decrease during a gout attack, so some time should elapse after the arthritis has resolved to check the level of uric acid in the blood. However, in a patient with gout, even with the treatment of arthritis, it is not necessary to expect a normal level of uric acid without antihyperuricemic therapy, that is, with normal values of this indicator, the presence of gout is doubtful. Of particular interest in the framework of the problem discussed is the study of inflammatory markers. On the one hand, uric acid is a “predictor” of cardiovascular events in the population, and its level is associated with the development of hypertension, insulin resistance and other classic cardiovascular risk factors. On the other hand, uric acid levels reflect the chronic inflammatory process in rheumatic diseases, including gout. X-ray examination of the joints. This helps to establish a differential diagnosis and identify typical signs of chronic gout. Radiologically diagnosed changes develop in 10-40% of cases after repeated attacks of arthritis and serve as one of the late manifestations of the disease. A characteristic radiological sign for late gout is the "shock" sign.

A number of authors have found that, as a rule, well-differentiated "perforated" periarticular erosions are detected radiographically 6-12 years after the first acute attack of gout. Ultrasound in the diagnosis of gout. Ultrasound examination of joints (UTT) in gout is a promising direction. A characteristic ultrasound symptom in gout is the "double contour" symptom [14]. The "double contour" sign is associated with the ability of crystals to accumulate on the articular surface, which is detected by ultrasound as an additional bright line parallel to the line of passage of the subchondral bone. Another pathological

finding in gout that can be detected by ultrasound is tophus. Detection of tophus structures facilitates early therapeutic measures. The determination of renal excretion of uric acid, in particular the ratio of uric acid to creatinine, can be used to identify hyperproducers and to decide whether to prescribe uricosuric drugs. Magnetic resonance imaging (MRI) is particularly useful in gout for the detection of tophus masses, since synovitis, synovial fluid, periarticular tissue damage, and edema are not characteristic of gout. Several authors have shown that tophus formation can occur early in the disease and even before the appearance of typical signs of arthritis. MRI can be informative in the differential diagnosis. Thus, the use of MRI in the differential diagnosis of tophus and neoplasms and infectious processes has been described. In addition, MRI can be a good monitoring tool during antihyperuricemic therapy due to its ability to assess the size and number of tophus over time. Computed tomography (CT) is another potentially useful method for imaging tophus. The method allows to detect tophus localized both in the bone and in the tendon and soft tissue area. So far, the method has not found widespread application, primarily due to its high cost. Thus, a review of the recent literature shows that at present the problem of early diagnosis and modern treatment of arterial hypertension in patients with gout and asymptomatic hyperuricemia remains open.

REFERENCES:

1. Nozima, S. (2023). CLINICAL AND PATHOGENETIC ASPECTS OF THE COURSE AND TREATMENT OF HYPERTENSION. *EUROPEAN JOURNAL OF MODERN MEDICINE AND PRACTICE*, 3(11), 25-29.
2. Erkinjonovna, S. N. (2024). THE RELATIONSHIP BETWEEN FOOD AND BLOOD PRESSURE. *EUROPEAN JOURNAL OF MODERN MEDICINE AND PRACTICE*, 4(4), 191-197.
3. Sharapova, N. (2023). ARTERIAL GIPERTENZIYA VA SEMIZLIK KASALLIKLARINING O'ZARO BOG'LIQLIK SABABLARI VA METABOLIK SINDROMLAR. *Центральноазиатский журнал образования и инноваций*, 2(11 Part 2), 174-179.

4. Шарапова, Н. (2023). КЕКСА ВА ҚАРИ ЁШЛИ АЁЛЛАРДА БЕЛ АЙЛАНАСИНИНГ ЖИСМОНИЙ ФАОЛЛИК БИЛАН БОҒЛИҚЛИГИ ҚИЁСИЙ ТАҲЛИЛИ. *Центральноазиатский журнал образования и инноваций*, 2(12 Part 2), 127-133.
5. Erkinjonovna, S. N. (2023). DIABETES MELLITUS IN PREGNANT WOMEN. *Best Journal of Innovation in Science, Research and Development*, 110-116.
6. Erkinjonovna, S. N. (2024). CHARACTERISTICS OF DENTAL PROSTHESES WEARING IN PATIENTS WITH TYPE 2 DIABETES ACCORDING TO KIDNEY IMPAIRMENT. *PEDAGOG*, 7(1), 84-88.
7. Erkinjonovna, S. N. (2024). THE BEST WAYS TO CONTROL HIGH BLOOD PRESSURE WITHOUT MEDICATION. *Journal of new century innovations*, 47(2), 175-183.
8. Sharapova, N. E. (2024). MODERN DIAGNOSTICS OF RESISTANT ARTERIAL HYPERTENSION. *EUROPEAN JOURNAL OF MODERN MEDICINE AND PRACTICE*, 4(11), 436-442.
9. Sharapova, N. E. (2024). Acute Bronchitis: Causes, Symptoms, and Treatment with Home Remedies. *American Journal of Bioscience and Clinical Integrity*, 1(10), 77-84.
10. Mavlonova, U. K., & Maxmurovna, M. K. (2020). The introduction of irony in english and uzbek literature. *International Engineering Journal For Research & Development*, 5(3), 4-4.
11. Абдуллаева, Л. С., Самадова, С. А., & Махмурова, М. (2014). Современные методы преподавания иностранных языков. Коммуникативный метод. *Наука. Мысль: электронный периодический журнал*, (6), 73-76.
12. Khamdamovna, M. U., & Khalimovna, M. M. (2020). ANALYSIS OF SITUATIONAL IRONY IN LITERATURE. *Academy*, (11 (62)), 26-27.
13. Mavlonova, U. K., Makhmurova, M. K., & Kodirov, T. D. (2020, June). Description of art media in literature. In LXIV International Correspondence

Scientific and Practical Conference “European Research: Innovation in science, education and technology» June (pp. 5-9).

14. Halimovna, M. M. (2020). The importance of a foreign language in higher education. *Вопросы науки и образования*, (8 (92)), 15-19.

15. Halimovna, M. M. (2020). LINGUISTIC BASIS OF TEACHING ORAL COMMUNICATION OF STUDENTS IN GERMAN LANGUAGE LESSONS. *BRIDGE TO SCIENCE: RESEARCH WORKS*, 65.

16. Tashpulatovna, B. V., Halimovna, M. M., & Tashpulatovich, B. M. (2021, November). Chet Tili O ‘Pgatishda O ‘Yin-Mashqlarning Roli. In " *ONLINE-CONFERENCES" PLATFORM* (pp. 139-144).

17. Halimovna, M. M. (2020). MODERN APPROACHES TO LEARNING A SECOND LANGUAGE. *Scientific reports of Bukhara State University*, 4(1), 182-185.

18. Khalimovna, M. M. (2021). The Role of Hadiths in Education and Upbringing. *CENTRAL ASIAN JOURNAL OF LITERATURE, PHILOSOPHY AND CULTURE*, 2(10).

19. Halimovna, M. M. (2021). The Role of Features of Linguocountrylearning in Teaching English. *Central Asian Journal of Literature, Philosophy and Culture*, 2(10), 64-68.

20. ERGASHEVA, G. T. (2024). OBESITY AND OVARIAN INSUFFICIENCY. *Valeology: International Journal of Medical Anthropology and Bioethics*, 2(09), 106-111.

21. Ergasheva, G. T. (2024). Modern Methods in the Diagnosis of Autoimmune Thyroiditis. *American Journal of Bioscience and Clinical Integrity*, 1(10), 43-50.

22. Tokhirovna, E. G. (2024). COEXISTENCE OF CARDIOVASCULAR DISEASES IN PATIENTS WITH TYPE 2 DIABETES. *TADQIQOTLAR. UZ*, 40(3), 55-62.

23. Musayeva Amina Karamatovna. (2024). Innovatsion Yondashuvlar Asosida Oliy Ta'lim Jarayoni Sifatini Oshirishning Metodologik Jihatlari. *Miasto Przyszłości*, 52, 59–64.
24. Karamatovna, M. A. . (2024). Important Aspects of Improving Students' Communicative Competence Today. *Miasto Przyszłości*, 55, 237–243.
25. Musayeva Amina Karamatovna. (2024). IMPROVING THE QUALITY OF THE HIGHER EDUCATION PROCESS BASED ON INNOVATIVE APPROACHES. *JOURNAL OF EDUCATION, ETHICS AND VALUE*, 3(9),95–100.
26. Toxirovna, E. G. (2024). DETERMINATION AND STUDY OF GLYCEMIA IN PATIENTS WITH TYPE 2 DIABETES MELLITUS WITH COMORBID DISEASES. *TADQIQOTLAR. UZ*, 40(3), 71-77.
27. Toxirovna, E. G. (2024). XOMILADORLIKDA QANDLI DIABET KELTIRIB CHIQUARUVCHI XAVF OMILLARINI ERTA ANIQLASH USULLARI. *TADQIQOTLAR. UZ*, 40(3), 63-70.
28. Toxirovna, E. G. (2024). QANDLI DIABET 2-TIP VA KOMORBID KASALLIKLARI BO'LGAN BEMORLARDA GLIKEMIK NAZORAT. *TADQIQOTLAR. UZ*, 40(3), 48-54.
29. Tokhirovna, E. G. (2024). MECHANISM OF ACTION OF METFORMIN (BIGUANIDE) IN TYPE 2 DIABETES. *JOURNAL OF HEALTHCARE AND LIFE-SCIENCE RESEARCH*, 3(5), 210-216.
30. Karamatovna, M. A. (2024). Integration of Distance and Full-Time Education as an Innovative Approach to Developing Communicative Competence. *EUROPEAN JOURNAL OF INNOVATION IN NONFORMAL EDUCATION*, 4(10), 5–13. Retrieved from <https://www.inovatus.es/index.php/ejine/article/view/4136>
31. Musayeva Amina Karamatovna. (2024). Innovatsion Yondashuvlar Asosida Oliy Ta'lim Jarayoni Sifatini Oshirishning Metodologik Jihatlari. *Miasto Przyszłości*, 52, 59–64. Retrieved from <https://miastoprzyszlosci.com.pl/index.php/mp/article/view/4535>

32. Karamatovna, M. A. (2024). Enhancing the Quality of Teaching in Higher Education: Strategies for Success. *American Journal of Science on Integration and Human Development* (2993-2750), 2(2), 97-103
33. Musaeva Amina Karamatovna. (2024). Formation of Students Communicative Competence. *Information Horizons: American Journal of Library and Information Science Innovation* (2993-2777), 2(1), 156–164.
34. Tokhirovna, E. G. (2024). THE ROLE OF METFORMIN (GLIFORMIN) IN THE TREATMENT OF PATIENTS WITH TYPE 2 DIABETES MELLITUS. *EUROPEAN JOURNAL OF MODERN MEDICINE AND PRACTICE*, 4(4), 171-177.
35. Эргашева, Г. Т. (2024). Эффект Применения Бигуанида При Сахарным Диабетом 2 Типа И Covid-19. *Research Journal of Trauma and Disability Studies*, 3(3), 55-61.
36. Тохировна, Е. Г. (2024). QANDLI DIABET 2 TUR VA YURAK QON TOMIR KASALLIKLARINING BEMOLARDA BIRGALIKDA KECISHI. *ОБРАЗОВАНИЕ НАУКА И ИННОВАЦИОННЫЕ ИДЕИ В МИРЕ*, 38(7), 202-209.
37. Эргашева, Г. Т. (2024). СОСУЩЕСТВОВАНИЕ ДИАБЕТА 2 ТИПА И СЕРДЕЧНО-СОСУДИСТЫХ ЗАБОЛЕВАНИЙ У ПАЦИЕНТОВ. *ОБРАЗОВАНИЕ НАУКА И ИННОВАЦИОННЫЕ ИДЕИ В МИРЕ*, 38(7), 219-226.
38. Эргашева, Г. Т. (2024). СНИЖЕНИЕ РИСКА ОСЛОЖНЕНИЙ У БОЛЬНЫХ САХАРНЫМ ДИАБЕТОМ 2 ТИПА И СЕРДЕЧНО-СОСУДИСТЫМИ ЗАБОЛЕВАНИЯМИ. *Образование Наука И Инновационные Идеи В Мире*, 38(7), 210-218.
39. Tokhirovna, E. G. (2024). CLINICAL AND MORPHOLOGICAL ASPECTS OF THE COURSE OF ARTERIAL HYPERTENSION. *Лучшие интеллектуальные исследования*, 12(4), 234-243.

40. Tokhirovna, E. G. Studying the Causes of the Relationship between Type 2 Diabetes and Obesity. *Published in International Journal of Trend in Scientific Research and Development (ijtsrd)*, ISSN, 2456-6470.
41. Toxirovna, E. G. (2024). ARTERIAL GIPERTENZIYA KURSINING KLINIK VA MORFOLOGIK JIHATLARI. *Лучшие интеллектуальные исследования*, 12(4), 244-253.
42. Эргашева, Г. Т. (2024). НОВЫЕ АСПЕКТЫ ТЕЧЕНИЕ АРТЕРИАЛЬНОЙ ГИПЕРТОНИИ У ВЗРОСЛОГО НАСЕЛЕНИЕ. *Лучшие интеллектуальные исследования*, 12(4), 224-233.
43. Эргашева, Г. Т. (2024). ФАКТОРЫ РИСКА РАЗВИТИЯ САХАРНОГО ДИАБЕТА 2 ТИПА. *ОБРАЗОВАНИЕ НАУКА И ИННОВАЦИОННЫЕ ИДЕИ В МИРЕ*, 36(5), 70-74.
44. Эргашева, Г. Т. (2024). ОСЛОЖНЕНИЯ САХАРНОГО ДИАБЕТА 2 ТИПА ХАРАКТЕРНЫ ДЛЯ КОГНИТИВНЫХ НАРУШЕНИЙ. *TADQIQOTLAR. UZ*, 30(3), 112-119.
45. Эргашева, Г. Т. (2023). Исследование Причин Связи Диабета 2 Типа И Ожирения. *Research Journal of Trauma and Disability Studies*, 2(12), 305-311.
46. Tokhirovna, E. G. (2024). Risk factors for developing type 2 diabetes mellitus. *ОБРАЗОВАНИЕ НАУКА И ИННОВАЦИОННЫЕ ИДЕИ В МИРЕ*, 36(5), 64-69.
47. Toxirovna, E. G. (2024). QANDLI DIABET 2-TUR VA O'LIMNI KELTIRIB CHIQRUVCHI SABABLAR. *Лучшие интеллектуальные исследования*, 14(4), 86-93.
48. Tokhirovna, E. G. (2023). Study of clinical characteristics of patients with type 2 diabetes mellitus in middle and old age. *Journal of Science in Medicine and Life*, 1(4), 16-19.
49. Toxirovna, E. G. (2024). GIPERPROLAKTINEMIYA KLINIK BELGILARI VA BERPUSHTLIKKA SABAB BO'LUVCHI OMILLAR. *Лучшие интеллектуальные исследования*, 14(4), 168-175.

50. Toxirovna, E. G. (2023). QANDLI DIABET 2-TUR VA SEMIZLIKNING O'ZARO BOG'LIQLIK SABABLARINI O'RGANISH. *Ta'lim innovatsiyasi va integratsiyasi*, 10(3), 168-173.
51. Makhmurova, M. H. (2024). Applications of the Latin Language in Speech. *American Journal of Alternative Education*, 1(8), 59-63.
52. Saloxiddinova, X. Y. (2024). Modern Views on the Effects of the Use of Cholecalciferol on the General Condition of the Bod. *JOURNAL OF HEALTHCARE AND LIFE-SCIENCE RESEARCH*, 3(5), 79-85.
53. Халимова, Ю. С., & Хафизова, М. Н. (2024). МОРФО-ФУНКЦИОНАЛЬНЫЕ И КЛИНИЧЕСКИЕ АСПЕКТЫ СТРОЕНИЯ И РАЗВИТИЯ ЯИЧНИКОВ (ОБЗОР ЛИТЕРАТУРЫ). *TADQIQOTLAR. UZ*, 40(5), 188-198.
54. Халимова, Ю. С. (2024). Морфологические Особенности Поражения Печени У Пациентов С Синдромом Мэллори-Вейса. *Journal of Science in Medicine and Life*, 2(6), 166-172.
55. Halimova, Y. S. (2024). Morphology of the Testes in the Detection of Infertility. *Journal of Science in Medicine and Life*, 2(6), 83-88.
56. Халимова, Ю. С., & Хафизова, М. Н. (2024). ОСОБЕННОСТИ СОЗРЕВАНИЕ И ФУНКЦИОНИРОВАНИЕ ЯИЧНИКОВ. *ОБРАЗОВАНИЕ НАУКА И ИННОВАЦИОННЫЕ ИДЕИ В МИРЕ*, 55(2), 188-194.
57. Хафизова, М. Н., & Халимова, Ю. С. (2024). МОТИВАЦИОННЫЕ МЕТОДЫ ПРИ ОБУЧЕНИИ ЛАТЫНИ И МЕДИЦИНСКОЙ ТЕРМИНОЛОГИИ. *ОБРАЗОВАНИЕ НАУКА И ИННОВАЦИОННЫЕ ИДЕИ В МИРЕ*, 55(2), 165-171.
58. Хафизова, М. Н., & Халимова, Ю. С. (2024). ИСПОЛЬЗОВАНИЕ ЧАСТОТНЫХ ОТРЕЗКОВ В НАИМЕНОВАНИЯХ ЛЕКАРСТВЕННЫХ ПРЕПАРАТОВ В ФАРМАЦЕВТИКЕ. *ОБРАЗОВАНИЕ НАУКА И ИННОВАЦИОННЫЕ ИДЕИ В МИРЕ*, 55(2), 172-178.
59. Saloxiddinova, X. Y., & Ne'matillaeva, X. M. (2024). FEATURES OF THE STRUCTURE OF THE REPRODUCTIVE ORGANS OF THE FEMALE

- BODY. *ОБРАЗОВАНИЕ НАУКА И ИННОВАЦИОННЫЕ ИДЕИ В МИРЕ*, 55(2), 179-183.
60. Халимова, Ю. С., & Хафизова, М. Н. (2024). КЛИНИЧЕСКИЕ АСПЕКТЫ ЛИЦ ЗЛОУПОТРЕБЛЯЮЩЕЕСЯ ЭНЕРГЕТИЧЕСКИМИ НАПИТКАМИ. *TADQIQOTLAR. UZ*, 40(5), 199-207.
61. Халимова, Ю. С., & Хафизова, М. Н. (2024). КЛИНИЧЕСКИЕ ОСОБЕННОСТИ ЗАБОЛЕВАНИЙ ВНУТРЕННИХ ОРГАНОВ У ЛИЦ, СТРАДАЮЩИХ АЛКОГОЛЬНОЙ ЗАВИСИМОСТЬЮ. *TADQIQOTLAR. UZ*, 40(5), 240-250.
62. Халимова, Ю. С., & Хафизова, М. Н. (2024). кафедра Клинических наук Азиатский международный университет Бухара, Узбекистан. *Modern education and development*, 10(1), 60-75.
63. Халимова, Ю. С., & Хафизова, М. Н. (2024). МОРФО-ФУНКЦИОНАЛЬНЫЕ И КЛИНИЧЕСКИЕ АСПЕКТЫ ФОРМИРОВАНИЯ КОЖНЫХ ПОКРОВОВ. *Modern education and development*, 10(1), 76-90.
64. Nematilloevna, K. M., & Salokhiddinovna, K. Y. (2024). IMPORTANT FEATURES IN THE FORMATION OF DEGREE OF COMPARISON OF ADJECTIVES IN LATIN. *ОБРАЗОВАНИЕ НАУКА И ИННОВАЦИОННЫЕ ИДЕИ В МИРЕ*, 55(2), 150-157.
65. KHALIMOVA, Y. S. (2024). MORPHOFUNCTIONAL CHARACTERISTICS OF TESTICULAR AND OVARIAN TISSUES OF ANIMALS IN THE AGE ASPECT. *Valeology: International Journal of Medical Anthropology and Bioethics*, 2(9), 100-105.
66. Salokhiddinovna, K. Y., Saifiloevich, S. B., Barnoevich, K. I., & Hikmatov, A. S. (2024). THE INCIDENCE OF AIDS, THE DEFINITION AND CAUSES OF THE DISEASE. *ОБРАЗОВАНИЕ НАУКА И ИННОВАЦИОННЫЕ ИДЕИ В МИРЕ*, 55(2), 195-205.