

## SOME MEDICINAL PROPERTIES AND ANALYSIS OF GLYCYRRHIZA GLABRA

*Researcher **Abdiravidov Temurbek Abdumannonovich**, 3<sup>rd</sup> year student of  
Clinical Pharmacy, Tashkent Pharmaceutical Institute*

*Scientific researcher, Tashkent Pharmaceutical Institute, 3<sup>rd</sup> year student of  
clinical pharmacology department, **Hikmatova Shukrona Ismatovna***

*The scientific supervisor is **Abzalova Zumrad Jurayevna**, assistant of the  
Department of Organic Synthesis of the Tashkent Pharmaceutical Institute*

**Keywords:** Antimicrobial, antioxidant, antiproliferative, Glycyrrhiza glabra, medicinal plants

### Abstract

Objective: Alternative medicine has an important place in the fight against many diseases in human history. The biological activity of Glycyrrhiza glabra L. was investigated in this study. Material and Method: The root parts of the plant were extracted with ethanol. Antioxidant and oxidant potentials were determined using Rel Assay kits. Antimicrobial activity was tested against standard bacteria and fungus strains using the agar dilution method. Antiproliferative activity was determined by MTT test against Lung Carcinoma Cell Line (A549). Result and Discussion: As a result of the studies, the TAS value of the plant was measured as  $8.770 \pm 0.171$ , TOS value as  $14.590 \pm 0.191$  and OSI value as  $0.167 \pm 0.005$ . Inhibition of the plant extract was observed against standard bacteria and fungus strains at ranging from 50-200  $\mu\text{g/mL}$  concentrations. In addition, it was determined that the plant extract displayed strong antiproliferative activity due to the increase in concentration. As a result of these studies, it has been determined that G. glabra can be used as an important natural antioxidant, antimicrobial and anticancer agent.

### INTRODUCTION

Many natural materials such as mushrooms, plants and animals are used in alternative medicine [1]. It occupies a very important place in the treatment of diseases due to the active substances produced by the plants. Many studies have shown that plants have anticancer, antioxidant, antimicrobial, antiproliferative, anti-inflammatory, DNA protective, antiallergic and hepatoprotective activities [2-4]. In our study, Glycyrrhiza glabra L. (Licorice) was used as a material. Licorice; It is a plant belonging to the genus Glycyrrhiza glabra, of the genus Glycyrrhiza of the Papilionaceae family. It is used for therapeutic purposes known in the history of ancient medicine in many civilizations such as Sumer, Mesopotamia, China, Greek

and Egypt in the world. Among the aromatic and medicinal plants, the licorice plant is popularly known by 14 different names such as "biyam, dye, piyam, sweet root, etc." [5]. Licorice plant in the world at 12, while in Turkey in 6 species of yellow-blue or brown color, is a perennial shrub plant ranging between 30-160 cm in length [6]. The roots of the licorice plant are biologically known as a source of magnesium and silicon. In addition, the active ingredient in the composition of the licorice plant, glycyrrhizin is 50 times sweeter than tea sugar and 150 times sweeter than sucrose. Glycyrrhiza acid, which is found in the sweetness of the roots, and its calcium and potassium salts, two of the substances such as sucrose and mannite come from [7, 8]. Therefore, as much more intense taste is obtained with less amount, it has been involved in the cuisine and food industry of many countries for centuries [9]. In addition, in the production of licorice honey, licorice sherbet, in the manufacture of tobacco, snuff and filter cigarettes, in the confectionery and beverage industry as a fragrance and flavoring, in cosmetics, velvet dyeing and shoe dyeing in the textile industry, making foam in fire fighting, in preparations prepared to kill insects, and in the food industry, there are areas of use such as adding fragrance to foods [10-12]. Licorice plant is widely used in food, confectionery, medicine and tobacco products as a flavoring agent known worldwide as "generally safe" (GRAS) [13]. In this study, antioxidant, oxidant, antimicrobial and antiproliferative activity of ethanol extract of root parts of *G. glabra* was determined.

## MATERIAL AND METHOD

### Laboratory Studies

Plant samples were collected from Duhok (Iraq). Soil and dust particles were removed from the root parts of the plant. It was then dried under suitable conditions. After drying the plant parts were pulverized and weighed 30 g. It was then extracted with ethanol for about 6 hours, for example at 50 °C. The solvents of the extracts were removed in a rotary evaporator and crude extracts were obtained.

### Antioxidant Parameters

The antioxidant and oxidant status of the plant extract was determined using Rel Assay TAS and TOS kits. TAS tests were performed according to the protocol specified in Erel [14] and Trolox was used as a calibrator. Results are shown in mmol Trolox equiv./L. TOS tests were performed according to the protocol specified in Erel [15] and hydrogen peroxide was used as a calibrator. Results are shown as  $\mu\text{mol H}_2\text{O}_2$  equiv./L. The oxidative stress index (OSI) was determined by proportioning the TOS value to the TAS value [16].

### Antimicrobial Activity Tests

The antimicrobial activity of the root parts of the plant against EtOH extract bacteria and fungus strains was determined using the agar dilution method [17-19].

The plant extract was adjusted with distilled water at 800-12.5  $\mu\text{g}/\text{mL}$  concentrations. Bacterial strains were set in Muller Hinton Broth medium. *Staphylococcus aureus* ATCC 29213, *S. aureus* MRSA ATCC 43300, *Enterococcus faecalis* ATCC 29212, *Escherichia coli* ATCC 25922, *Pseudomonas aeruginosa* ATCC 27853 and *Acinetobacter baumannii* ATCC 19606 were used as bacterial strains. Fungus strains were pre-cultured in RPMI 1640 Broth medium. *Candida albicans* ATCC 10231, *C. krusei* ATCC 34135 and *C. glabrata* ATCC 90030 were used as fungus strains. The extract concentration that inhibits the growth of bacteria and fungus strains was determined as the MIC value. Results were expressed in  $\mu\text{g}/\text{mL}$  [20-22].

### Antiproliferative Activity Tests

The antiproliferative activity of the EtOH extract of the plant was determined by MTT test on A549 lung cancer cells. Cells were separated after 70-80% confluence using 3.0 mL of Trypsin-EDTA solution (Sigma-Aldrich, MO, USA). It was then planted on plates and incubated for 24 hours. The plant extract (25, 50, 100, 200  $\mu\text{g}/\text{mL}$ ) was then adjusted at different concentrations. After the incubation period, the supernatants were dissolved in growth medium and replaced with 1 mg/mL MTT (Sigma). It was then incubated at 37 °C until a purple precipitate formed. The supernatants were then removed and dissolved by adding dimethyl sulfoxide (DMSO) (Sigma-Aldrich, MO, USA) to MTT absorbed by cells. Subsequently, plates were read at 570 nm using an Epoch spectrophotometer (BioTek Instruments, Winooska, VT) [23].

## RESULT AND DISCUSSION

**Antioxidant Activity** In recent years, it is known that the basis of many diseases are due to oxidative stress caused by reactive oxygen species. Oxidative stress is due to the disproportion between the formation and neutralization of prooxidants [24]. The level of endogenous oxidant compounds resulting from environmental factors as a result of metabolic activities is highly toxic when they accumulate in organisms. The antioxidant defense system plays a role in reducing the effects of oxidant compounds. If the antioxidant defense system is insufficient, oxidative stress occurs. In such cases, supplemental antioxidants are important in reducing the effect of oxidative stress. In this context, it is very important to identify new natural antioxidant agents [25]

In previous studies of *G. glabra*, it was reported that aqueous, methanol, ethanol extracts have antioxidant potential using different methods (Inhibition of b-carotene–linoleate bleaching, Hypochlorous acid-scavenging activity, Inhibition of myeloperoxidase-chlorinating system, Nitric oxide radical scavenging activity, Superoxide anion scavenging activity, Hydroxyl radical scavenging activity, DPPH radical scavenging activity, ABTS•+ cation Radical Scavenging, Fe<sup>+2</sup>/ascorbate

induced lipid peroxidation assay, Reducing power) [26-29]. In our study, TAS, TOS and OSI values were determined for the first time by using Rel Assay kits of *G. glabra*. As a result of the studies, it has been determined that *G. glabra* has an important antioxidant activity. In addition, TAS values of *R. coriaria* var. *zebaria*, *M. longifolia* subsp. *longifolia*, *A. calocephalum*, *S. papposa*, *F. platycarpa*, *T. spicata*, *G. tournefortii*, *R. crispus* and *A. millefolium* reported in the literature were reported as 7.342, 3.628, 5.853, 5.314, 5.688, 8.399, 6.831, 6.758 and 2.436 mmol/L, respectively. TOS values were reported as 5.170, 4.046, 16.288, 24.199, 15.552, 6.530, 3.712, 5.802 and 2.839  $\mu\text{mol/L}$ , respectively. OSI values were reported as 0.071, 0.112, 0.278, 0.456, 0.273, 0.078, 0.054, 0.086 and 0.083, respectively [30-38]. Compared to these studies, the TAS value of *G. glabra* was determined to be higher than *R. coriaria* var. *zebaria*, *M. longifolia* subsp. *longifolia*, *A. calocephalum*, *S. papposa*, *F. platycarpa*, *T. spicata*, *G. tournefortii*, *R. crispus* and *A. millefolium*. TAS value shows all of the antioxidant compounds produced in the plant [30]. As seen in our study, it has been determined that *G. glabra* has a very important antioxidant potential.

When TOS values were examined, it was determined that *G. glabra* was lower than *S. papposa* and *F. platycarpa*, and higher than *Rhus coriaria* var. *zebaria*, *Mentha longifolia* subsp. *longifolia*, *A. calocephalum*, *T. spicata*, *G. tournefortii*, *R. crispus* and *A. millefolium*. The TOS value indicates all of the oxidant compounds produced by the environmental effects in the plant [30]. It is seen that the oxidant levels of the plant used in our study are at normal levels. When OSI values were examined, it was determined that *G. glabra* was lower than *A. calocephalum*, *S. papposa* and *F. platycarpa*, and higher than *R. coriaria* var. *zebaria*, *M. longifolia* subsp. *longifolia*, *T. spicata*, *G. tournefortii*, *R. crispus* and *A. millefolium*. The OSI value shows how much oxidant compounds produced in the plants are suppressed by the antioxidant defense system. A low OSI value indicates that the antioxidant defense system of the plant works well [30]. In our study, it was determined that the antioxidant defense system of *G. glabra* was sufficient in suppressing oxidant compounds. As a result, it was determined that *G. glabra* has significant antioxidant activity.

### Antimicrobial Activity

Today, many diseases occur due to microorganisms. Antibiotics are used extensively in the treatment of microorganism-based diseases. Today, resistant strains are emerging due to the unconscious use of antibiotics [39]. Antibiotics used against resistant microorganisms are insufficient. In addition, due to the possible side effects of chemical antibiotics, the tendency towards natural products is increasing. In this context, the discovery of new antimicrobial drugs is inevitable [40]. In this study, the activities of *G. glabra* against bacteria and fungi were investigated.

In previous studies, it has been reported that methanol extracts of *G. glabra* are effective against *Staphylococcus aureus*, *Bacillus megaterium*, *B. subtilis*, *Sarcina lutea*, *Salmonella paratyphi*, *S. typhi*, *Escherichia coli*, *Shigella dysenteriae*, *Vibrio minicus*, *V. parahemolyticus* and *Pseudomonas aeruginosa* at different concentrations [41]. In a different study, ethanol extracts of *G. glabra* were reported to be effective against *Candida albicans*, *Staphylococcus aureus*, *Bacillus subtilis*, *Enterococcus faecalis*, *Escherichia coli*, *Pseudomonas aeruginosa* and *Klebsiella pneumoniae* [42]. Ethanolic, hexane fraction, ethyl acetate fraction and methanol fraction of *G. glabra* have been reported to be effective against *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Streptococcus mutans*, *Bacillus subtilis*, *Enterococcus faecalis*, *Klebsiella pneumoniae*, *Salmonella typhi*, *Yersinia enterocolitica*, *Enterobacter aerogens* and *Escherichia coli* [43]. In our study, *G. glabra* was determined to be effective against *A. baumannii*, *C. glabrata* and *C. albicans* at 25 µg/mL, *S. aureus*, *S. aureus* MRSA and *C. krusei* at 50 µg/mL, *E. faecalis* and *P. aeruginosa* at 100 µg/mL, *E. coli* at 200 µg/mL extract concentrations. As a result, it was determined that *G. glabra* has antibacterial and antifungal activities.

### Conclusion

In this study, some medicinal properties of the root parts of *G. glabra* were determined. As a result of the studies, it was determined that the root extracts of the plant exhibit significant antioxidant, antimicrobial and antiproliferative activity. As a result, it is thought that *G. glabra* can be used as a natural material in pharmacological designs.

### REFERENCES

1. Korkmaz, A.I., Akgul, H., Sevindik, M., Selamoglu, Z. (2018). Study on determination of bioactive potentials of certain lichens. *Acta Alimentaria*, 47(1), 80-87. [CrossRef]
2. Ben-Shabat, S., Yarmolinsky, L., Porat, D., Dahan, A. (2020). Antiviral effect of phytochemicals from medicinal plants: applications and drug delivery strategies. *Drug Delivery and Translational Research*, 10(2), 354-367. [CrossRef]
3. Li, Y., Kong, D., Fu, Y., Sussman, M.R., Wu, H. (2020). The effect of developmental and environmental factors on secondary metabolites in medicinal plants. *Plant Physiology and Biochemistry*, 148, 80-89. [CrossRef]
4. Mohammed, F.S., Pehlivan, M., Sevindik, E., Akgul, H., Sevindik, M., Bozgeyik, I., Yumrutas, O. (2021). Pharmacological properties of edible *Asparagus acutifolius* and *Asparagus officinalis* collected from North Iraq and Turkey (Hatay). *Acta Alimentaria*, 50(1), 136-143. [CrossRef]

5. Pastorino, G., Cornara, L., Soares, S., Rodrigues, F., Oliveira, M.B.P. (2018). Liquorice (*Glycyrrhiza glabra*): A phytochemical and pharmacological review. *Phytotherapy research*, 32(12), 2323-2339. [CrossRef]
6. Kılınç, F., Demircan, F., Yıldırım, Y., Yılmaz, Z., Pekkolay, Z., Tuzcu, A.K. (2014). Meyan Kökü (Likoris) Kullanımına Bağlı Hipertansiyon ve Hipokalemi ile Seyreden 5 Vakanın Değerlendirilmesi. *Sakarya Tıp Dergisi*, 4, 186-190. [CrossRef]
7. Nazari, S., Rameshrad, M., Hosseinzadeh, H. (2017). Toxicological effects of *Glycyrrhiza glabra* (licorice): a review. *Phytotherapy research*, 31(11), 1635-1650. [CrossRef]
8. Alagawany, M., Elnesr, S.S., Farag, M.R., El-Hack, A., Mohamed, E., Khafaga, A.F., Taha, A.E., Tiwari, R., Yattoo, M.I., Bhatt, P., Marappan, G., Dhama, K. (2019). Use of licorice (*Glycyrrhiza glabra*) herb as a feed additive in poultry: Current knowledge and prospects. *Animals*, 9(8), 536. [CrossRef]
9. Cuvelier, M. E., Berset, C., Richard, H. (1994). Antioxidant Constituents in Sage (*Salvia officinalis*). *Journal of Agricultural and food chemistry*, 42, 665-669. [CrossRef]
10. Kikuzaki, H., Nakatani, N. (1993). Antioxidant Effects of Some ginger constituents. *Journal of food science*, 58, 1407-1410. [CrossRef]
11. Shahidi, F. (2000). Antioxidants in Food and Food Antioxidants. *Food/nahrung*, 44, 158-163. [CrossRef]
12. Dang, M.N., Takácsová, M., Nguyen, D.V., Kristiánová, K. (2001). Antioxidant Activity of Essential Oils from Various Spices. *Food/nahrung*, 45, 64-66. [CrossRef]